

Yu-Feng Wang, Todd A. Ponzio and Glenn I. Hatton

Am J Physiol Regulatory Integrative Comp Physiol 290:1191-1198, 2006. First published Dec 1, 2005;
doi:10.1152/ajpregu.00725.2005

You might find this additional information useful...

This article cites 26 articles, 13 of which you can access free at:

<http://ajpregu.physiology.org/cgi/content/full/290/5/R1191#BIBL>

This article has been cited by 1 other HighWire hosted article:

Dominant Role of $\beta\gamma$ Subunits of G-Proteins in Oxytocin-Evoked Burst Firing

Y.-F. Wang and G. I. Hatton
J. Neurosci., February 21, 2007; 27 (8): 1902-1912.
[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Updated information and services including high-resolution figures, can be found at:

<http://ajpregu.physiology.org/cgi/content/full/290/5/R1191>

Additional material and information about *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology* can be found at:

<http://www.the-aps.org/publications/ajpregu>

This information is current as of November 24, 2009 .

The American Journal of Physiology - Regulatory, Integrative and Comparative Physiology publishes original investigations that illuminate normal or abnormal regulation and integration of physiological mechanisms at all levels of biological organization, ranging from molecules to humans, including clinical investigations. It is published 12 times a year (monthly) by the American Physiological Society, 9650 Rockville Pike, Bethesda MD 20814-3991. Copyright © 2005 by the American Physiological Society. ISSN: 0363-6119, ESN: 1522-1490. Visit our website at <http://www.the-aps.org/>.

CALL FOR PAPERS | *Neurohypophyseal Hormones: From Genomics and Physiology to Disease*

Autofeedback effects of progressively rising oxytocin concentrations on supraoptic oxytocin neuronal activity in slices from lactating rats

Yu-Feng Wang, Todd A. Ponzio, and Glenn I. Hatton

Department of Cell Biology and Neuroscience, University of California, Riverside, California

Submitted 12 October 2005; accepted in final form 22 November 2005

Wang, Yu-Feng, Todd A. Ponzio, and Glenn I. Hatton. Autofeedback effects of progressively rising oxytocin concentrations on supraoptic oxytocin neuronal activity in slices from lactating rats. *Am J Physiol Regul Integr Comp Physiol* 290: R1191–R1198, 2006. First published December 1, 2005; doi:10.1152/ajpregu.00725.2005.—Suckling stimuli induce somatodendritic oxytocin (OT) release from supraoptic nucleus (SON) neurons, which raises intranuclear OT concentrations and contributes to the effectiveness of the milk-ejection reflex. To clarify how such changes in OT concentrations modulate the activity of OT neurons, we examined OT effects using whole cell patch-clamp recordings from SON neurons in slices from lactating rats. Progressive increases from extremely low OT concentrations (0.1–10 fM) to high concentrations (0.1–10 nM) induced excitation and subsequent spike frequency reduction (SFR) in OT neurons. Significant effects of OT on firing rates were observed starting at 1 fM, reached peak level from 1 fM to 1 pM before SFR occurred in most neurons. The buildup of OT concentrations progressively promoted depolarization of membrane potential, spike broadening, decreases in spike amplitude, and increases in the rise time of spike afterhyperpolarizations, which were unrelated to firing rate. However, intermittent application of OT (1 fM, 1 pM, and 1 nM, each for 5 min) evoked dose-dependent excitation but not the SFR. Application of 1 pM OT for 40 min simulated the effects of progressively increasing OT concentrations. Vasopressin neurons were also activated by OT but did not show SFR. Consistent with presynaptic loci of OT action, ionotropic glutamate receptor antagonists reduced OT effects on firing rate, whereas bicuculline did not change the excitatory effects. These results suggest that the specific autoregulatory effects of OT, and perhaps other neuropeptides as well, are time and concentration dependent.

integration; lactation; neuropeptide; spike frequency reduction; synaptic transmission

THE MAGNOCELLULAR HYPOTHALAMO-neurohypophyseal system, of which oxytocin (OT) secretion is an integral functional part, serves as a model for neurosecretion. During suckling, extracellular OT concentrations in the maternal supraoptic nucleus (SON) increase significantly (20, 21). This facilitates activation of OT neurons (27). Intracerebroventricular injection of OT also facilitates the activation of OT neurons, while injection of OT antagonists inhibits ongoing bursts of OT neurons (10, 20). Myriad *in vitro* studies revealed that OT induced its own release (19), resulting in excitation of OT neurons (4, 28). Moreover, OT reduces the amplitude of inhibitory postsynaptic

currents (IPSCs) (3). In contrast to this evidence supporting its facilitatory effects, OT has also been found to reduce the amplitude of evoked postsynaptic currents (EPSCs) at presynaptic terminals on SON neurons (14), which possibly reduces excitatory drive. These results raise questions about the integrative actions of OT. As previous studies are mostly based on brief applications of single doses of OT, they did not address questions of the effects of gradual increases in OT concentrations, as occur *in vivo*, over longer periods of suckling. Because these increasing levels of OT may cause secondary and hitherto unknown responses in OT neurons, it was important to determine the effects elicited by slow, progressive increases in OT concentration.

To acquire these data relating to the mechanisms underlying OT autoregulation, we examined the effects of gradually increasing OT concentrations on the electrical activity of OT neurons using whole cell patch-clamp recordings in brain slices from lactating rats. In pilot experiments, in which extremely low OT concentrations were tested, we determined that OT *in vitro* could exert excitatory effects at concentrations that were much lower than the previously measured physiological range. We found that OT did cause decreased responsiveness, akin to desensitization, after initial excitatory effects. These actions are achieved via OT's actions at both local neural circuits and on membrane electrical characteristics.

MATERIALS AND METHODS

All procedures in the animal experiments were in accordance with the guidelines on the use and care of laboratory animals set by National Institutes of Health (NIH) and approved by the Institutional Animal Care and Use Committee of the University of California, Riverside.

Electrophysiology. Sprague-Dawley (Holtzman strain) rats lactating for 8–13 days were used for the experiments. Rats were decapitated with a guillotine. Brains were quickly removed and put in oxygenated, ice-cold artificial cerebrospinal fluid (aCSF) for 1 min. The aCSF contained (in mM): 126 NaCl, 2.5 KCl, 1.3 MgSO₄, 2.4 CaCl₂, 1.3 NaH₂PO₄, 26 NaHCO₃, 10 glucose, 0.2 ascorbic acid, and pH 7.4 adjusted with MOPS (~2 mM). The osmolality was adjusted to 300 mosmol/kgH₂O. The aCSF was filtered (0.22 μm) and maintained with 95% O₂-5% CO₂ gas mixture. Hypothalami were dissected from the brain and cut coronally into cold medium on a Vibratome into 300-μm-thick slices. After preincubation at room

Address for reprint requests and other correspondence: Yu-Feng Wang, Dept. of Cell Biology and Neuroscience, Univ. of California, Riverside, Riverside, CA 92521 (E-mail: yufengw@ucr.edu).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

temperature ($22 \pm 1^\circ\text{C}$) for at least 1 h (usually more than 6 h), the slices were placed in a perfusion chamber and used for electrical recordings at 35°C . Whole cell patch-clamp methods were used to record membrane potential (E_m) and action potentials. Patch pipette filling solution contained (in mM): 145 K-gluconate, 10 KCl, 1 MgCl_2 , 10 HEPES, 1 EGTA, 0.01 CaCl_2 , 2 Mg-ATP, 0.5 $\text{Na}_2\text{-GTP}$, and pH 7.3, adjusted with KOH. For immunocytochemical identification of recorded neurons, 0.05% Lucifer yellow (LY) was added to the pipette solution. Patch electrodes were visually guided onto SON cells via an upright microscope (Leica DM LFS) equipped with water immersion objectives, infrared/differential interference contrast, and filters for fluorescent microscopy. Whole cell recordings were obtained from the somata of SON magnocellular neurons during perfusion of aCSF via a gravity-fed perfusion system at a rate of 1.2–1.5 ml/min. An Axoclamp 2B amplifier was used for collecting electrical signals that were filtered and sampled at 5 kHz by Clampex 9 software through a 1320 AD/DA converter (Axon Instruments). Data were stored in a PC computer for offline analysis.

Immunocytochemical identification of OT neurons. After recording with electrodes containing LY in the pipette solution, slices were fixed overnight with 4% paraformaldehyde at 4°C and treated with 0.3% Triton X-100 for 30 min. After incubation with mouse monoclonal OT-neurophysin antibody (PS38, 1:400 dilution) for 4 h at room temperature, a goat anti-mouse antibody (Alexa Fluor 647 labeled, 1:1,000) was applied for 1.5 h to label OT neurons. Sections were sealed on glass slides with Vectashield to avoid bleaching and examined with a laser scanning confocal microscope (Leica TCP SP2) in sequential scanning mode. Three-dimensional overlap of LY images with PS38/Alexa Fluor 647 immunostaining was taken as an identification of OT neurons.

Agents. 6-cyano-7-nitroquinoxaline-2, 3-dione (CNQX), bicuculline methiodide, LY (K^+ salt), and OT were obtained from Sigma (St. Louis, MO). (+)-MK-801 hydrogen maleate (MK 801) was from Tocris (Ellisville, MO). The monoclonal antibody against OT-neurophysin (PS38) was kindly provided by Dr. H. Gainer (NIH, Bethesda, MD). Alexa Fluor 647-labeled goat anti-mouse IgG was from Molecular Probes (Eugene, OR). Vectashield was purchased from Vector Laboratories (Burlingame, CA).

Data collection and analysis. Neurons that fired continuously and showed sustained outward rectification (SOR) in response to 11 steps of hyperpolarizing pulses (each lasting 1,200 ms) (25) were taken as putative OT neurons. Putative vasopressin (VP) neurons were those that fired phasically and showed clear plateau potentials. Exemplary neurons in each group and neurons that could not be classified by electrophysiological criteria were further identified subsequently by immunocytochemistry, and silent neurons were not analyzed in this study. Spikes and spike afterhyperpolarizations (AHPs) in patch-clamp recordings were analyzed with Clampfit 9 software (Axon Instruments) after filtering at 1 Hz. ANOVA, Student's *t*-tests and χ^2 tests were used for statistical analyses, and $P < 0.05$ was considered significant. Liquid junction potentials (-8 to -11 mV) were corrected according to the actual value measured in different groups of pipettes. All measures were expressed as means \pm SE, except as otherwise indicated in the results.

RESULTS

Recordings were made from 54 neurons in the SON in vitro using patch-clamp methods. Forty-five of these showed distinctly positive SOR responses and were electrically classified as OT neurons; six were phasic, VP neurons, and three neither showed clear SOR responses nor were phasic. Eleven neurons, including the three that could not be classified by electrophysiological criteria, were further identified immunocytochemically as being OTergic. The six VP neurons were used for controls.

Effects of progressively increasing OT concentrations on firing activity of OT neurons. It should be noted at the outset that the neurons recorded in this configuration tend to be superficial in the slice and, thus, well perfused by medium. This likely keeps concentrations of endogenously released substances from accumulating, as they might at sites deeper in the slice. OT is a crucial neuroactive substance in the SON during milk ejection (27). To examine the effects of OT on the electrical activity of OT neurons, we simulated the physiological changes in OT by bath application in progressively increasing concentrations (0.1 fM–10 nM in nine steps, each for 5 min). OT exerted an excitatory effect on nearly all OT neurons tested by increasing the firing rate by more than 20% of control level in a 5-min period (Fig. 1A). This excitatory effect started at the low concentrations (8/9 at 1 fM), then became less excitatory, and even reversed to silence at higher concentrations in all nine neurons, appearing to cause spike frequency reduction (SFR). The firing rate changes (excitation and ensuing SFR) through the nine doses were statistically significant ($P < 0.01$ by ANOVA) after square root transformation of the original data. Accompanying the firing rate changes, E_m depolarized significantly, which appeared to be time and dose dependent.

Dose- and time-related effects of OT on electrical activity of OT neurons. The effect of progressively increasing OT concentrations may reflect both time and dose in OT actions. We first tested the dose effects by applying OT in three concentrations (1 fM, 1 pM, and 1 nM, each for 5 min, interrupted by 10-min washout). Different from the effect of progressively increasing concentrations, the intermittently applied OT caused dose-dependent increases in firing rate without SFR (Fig. 1B). This result indicates that a recovery time to restore the receptor to its prebound state appears to be necessary for restored activation of OT neurons and is supported by the next set of experiments. Prolonged application of a single OT dose (1 pM, 40 min, $n = 5$) did cause initial excitation (4/5) and ensuing SFR (Fig. 1C). Thus the effects of progressively increasing OT concentrations on the firing rate are both time and dose dependent.

Corresponding to the responses from OT neurons, the progressively increasing OT concentrations also caused excitation of phasically firing, putative VP SON neurons (for examples of phasically firing cells recorded in vitro, see Refs. 11 and 12). In contrast to the effects on OT neurons, the excitation in VP neurons occurred with longer latency and only at higher OT concentrations: the initial excitation appeared during 1–10 pM range (5/6), and was not followed by SFR in four of the six neurons recorded (data not shown).

Effects of progressively increasing OT concentrations on spike and other membrane electrical characteristics. The firing rates of OT neurons are also influenced by other membrane electrical activity (e.g., spiking threshold, E_m level, spike AHPs, and others). Therefore, we further analyzed the effect of dynamic changes in OT concentrations on these electrical activities. Besides evoking the dual firing rate changes and E_m depolarization, OT also caused a reduction in spike amplitude, increase in spike duration, prolongation of spike rise and decay time courses, elevation of spike threshold (from -61.0 ± 3.2 to -54.6 ± 2.6 mV at 1 pM, $n = 9$, $P < 0.05$), and an increase in rise time of the spike AHP (Fig. 2A). Despite dramatic effects of OT on other electrical properties of OT neurons,




Fig. 1. Progressively increasing oxytocin (OT) concentrations induced excitation and subsequent depression in firing activity of OT neurons. A–C: voltage recordings of OT neuronal activity at resting membrane potential. A: effect of progressively increasing OT concentrations (from 0.1 fM to 10 nM, 5 min for each dose) on firing activity of an OT neuron (A1). At left is the initial membrane potential (E_m), corrected for liquid junction potential. Washout partially reinstated control levels of E_m and firing. The short solid lines above the recordings in A1, B1, and C1 indicate control levels of spike amplitude. The long dotted line below the traces corresponds to the resting E_m in the first trace. Note, after the initial excitation, the neuron became less active, accompanied by dramatic E_m depolarization and obvious spike amplitude reduction before becoming silent at 10 nM OT concentration. These responses were typical of recorded neurons. A2, left: scattergram of frequency (basal firing rate, FR, vs. FR during OT application). A2, middle and right: firing rate (Freq.) and resting E_m (RMP), respectively. B: three OT concentrations applied with interapplication intervals of 10 min (1 fM, 1 pM, and 1 nM, each for 5 min). B1: example of one neuron. B2: summary graphs. C: effects of prolonged single OT concentration (1 pM, 40 min) on the activity of OT neurons. C1: example of one neuron. C2: summary graphs. * $P < 0.05$; ** $P < 0.01$ compared with control by paired Student's t -test after ANOVA evaluation; n = the number of neurons.

membrane conductance ($n = 9$) did not change significantly. Different from the effects on firing rate, the effects of OT on the above parameters were dose and time dependent, suggesting different underlying regulatory mechanisms. However, increases in the number of spike clusters were similar to effects of OT on the firing rate. Time course of the subthreshold E_m

depolarization (STD) was extended significantly around 1 pM OT, then returned to basal level at higher concentrations (Fig. 2A₂).

It is noteworthy that the beginning of recovery (usually within 10 min) was significantly faster for firing rates than for spike amplitudes, spike width, E_m , AHP amplitude, or rise and




Fig. 2. Membrane electrical characteristics of OT neurons are changed by progressive increases in OT concentration. *A*: voltage recordings of OT neuronal activity. *A1*: example of one neuron. Dashed arrows in *A2* show the origins of individual spikes expanded in *A2* and in *A3*. OT progressively depolarized the E_m while inducing spike frequency reduction (SFR) after a short increase before 1 pM application. *A2*: dotted lines below selected traces indicate the time courses of STD leading to spike generation. *A3*: individual examples showing spike details. The long dashed line corresponds to spike threshold. Distance between short horizontal dashed lines and the long horizontal line indicates the amplitude of spike afterhyperpolarization (AHP). Distances between vertical lines (*A3*, left and middle) below and above long horizontal line indicate the 50% amplitude (Amp.) width of the AHPs and 50% amplitude width of spikes (between arrows), respectively. *B*: summary graphs showing the effects of OT on the amplitude and width of the spike, and their rising (RT) and decay (DT) time constants, the STD, and incidence of spike clusters (three consecutive spikes with interspike intervals of less than 100 ms) besides the RT of the AHPs. Control values for the Amp, Width, RT, DT, STD, spike clusters, and AHP-RT are 62.1 mV, 1.7 ms, 0.5 ms, 4.8 ms, 45 ms, 0.9 cluster/5 min, 1.7 ms, respectively. * $P < 0.05$; ** $P < 0.01$ compared with control. Other annotations are the same as in Fig. 1.

decay time courses of AHPs. This provides additional evidence for the existence of different regulatory processes involved in firing rate vs. most of the other electrical characteristics (summarized in Fig. 2*B*).

Fast synaptic inputs and OT effects. The autoregulation of OT neuronal activity may result from both direct and indirect effects of OT by modulation of other neuroactive substances, that is, those coexisting with OT in the extracellular environment (7, 27). Because glutamate and GABA inputs are the main sources of direct synaptic activity on OT neurons and are known to be modulated by OT, we examined the effects of blocking fast glutamatergic and GABAergic synaptic transmission on OT actions. Application of 10 μ M CNQX, 20 μ M MK 801, and 20 μ M bicuculline produced a brief increase in firing activity, then silence in 7 of 9 neurons, during which time the E_m depolarized by 3–10 mV (6.5 ± 1.7 mV). The presence of these blockers reduced and delayed the excitatory effect of OT. Three neurons excited by OT showed no SFR. These results suggest that the excitatory effects of OT and the SFR are closely related to synaptic inputs, besides confirming a dominant GABAergic innervation of OT neurons in the SON.

To distinguish these presynaptic actions, we further examined effects of OT after blocking ionotropic glutamate recep-

tors. In the presence of 10 μ M CNQX and 20 μ M MK 801 (Fig. 3*A*), spontaneous firing activity was reduced in a majority of OT neurons (6/9), addition of OT did not significantly increase the excitability in general, and did not cause SFR. However, other effects exerted by OT in normal aCSF were maintained, for example, E_m depolarization (from -60.7 ± 2.4 to -57.6 ± 2.1 mV at 1 pM, $n = 9$, $P < 0.05$), spike amplitude reduction, and spike broadening were observed, although they were not as dramatic as those seen in the normal aCSF (Fig. 3*A2* and 3*B*). The effects on spike amplitude and spike width were blocked by intracellularly loading 2 mM GDP β S, an inhibitor of G-proteins (Fig. 3*C*).

By contrast, the presence of bicuculline (20 μ M) to block GABA_A-mediated synaptic transmission ($n = 9$), depolarized E_m in all neurons (>3 mV in 6/9) and increased firing rates in most of them (Fig. 4). Addition of OT caused qualitatively similar changes in the membrane electrical features, to those seen in the absence of bicuculline. However, the SFR phenomena were still present during GABA_A receptor blockade. These results suggest that the excitatory effects of OT are associated with increased glutamate release and decreased GABA release and that the SFR phenomena are related to increased excitatory actions rather than to reduced inhibitory synaptic input.




Fig. 3. Blocking ionotropic glutamate receptors reduced OT excitatory effects. A–C: influence of blocking fast synaptic transmission on the effects of OT on the firing activities. A: progressively increasing OT concentrations failed to evoke SFR in the presence of 10 μ M 6-cyano-7-nitroquinoxaline-2, 3-dione (CNQX) and 20 μ M MK801 (CM), but still caused significant effects on spike amplitude and duration. B: Summary graphs (cf. Fig. 2). Control values for the Freq, Amp, Width, RT, DT, STD, spike clusters, and AHP-RT are 0.3 Hz, 59.1 mV, 1.5 ms, 0.6 ms, 35.8 ms, 58.9 ms, 0.15 clusters/5 min, and 0.9 ms, respectively. C: effect of blocking G-protein mediation by including 2 mM GDP- β -S in the pipette solution, on OT actions in the presence of CNQX and MK801 ($n = 3$). OT effects on the E_m , spike amplitude, and duration were further reduced. Other annotations are the same as in Figs. 1 and 2.

DISCUSSION

Progressively increasing OT concentrations caused excitation and SFR, as well as a series of membrane electrical changes. Glutamatergic inputs promoted the excitatory action of OT and the subsequent SFR, while GABA played antagonistic roles in excitatory response of OT neurons. In addition, postsynaptic actions of OT determined most other electrical responses.

Excitatory effect of OT on the firing activity of OT neurons. High sensitivity of neurons to neuropeptides is commonly observed; however, few if any electrophysiological studies

have reported femtomolar effects of neuropeptides. In the present work, OT triggered or increased firing activity of OT neurons, starting at 1 fM concentration. These are even lower than basal OT concentrations in the SON estimated by microdialysis (16, 21, 22). The nature of the perfusion of the tissue in vitro studies may remove endogenous OT from the surface of the slices, allowing higher accessibility and sensitivity to OT than the neurons recorded at sites deeper in the slice in extracellular or intracellular recordings. This, together with the relatively long time of OT application (providing enough time




Fig. 4. Blocking GABA_A receptors did not dramatically change the actions of OT. *A*: example showing the influence of blocking GABA_A receptor with 20 μM bicuculline (Bic) on the effects of OT. *B*: summary graphs (cf. Figs. 2 and 3). Control values for the Freq, Amp, Width, RT, DT, STD, spike clusters, and AHP-RT are 2.3 Hz, 63 mV, 2.5 ms, 1.2 ms, 109 ms, 38 ms, 1.1 clusters/5 min, and 1.2 ms, respectively. Other annotations are the same as in Figs. 1 and 2.

for temporal summation), using normal aCSF in observations but not low Ca²⁺ to elevate neuronal excitability artificially, carefully avoiding remaining effects of high-dose OT by thoroughly cleaning the perfusion system between tests, may account for the biological effect of lower OT concentration on OT neurons. It has been reported that 0.2 nM OT could excite hypothalamic neurons (15); and the natural cellular environment (the presence of Mg²⁺ and cholesterol) may also be a favorable condition for the functioning of saturable high-affinity OT receptors (13). Finally, what we measured was the electrophysiological output of the OT receptor in neurons, which may have different characteristics (affinities, K_Ds, etc.) from those in peripheral tissue where these measurements have previously been made. Thus it is reasonable to find the effect of this low concentration of OT in cellular function.

Because OT can activate both OT receptors and V₁ receptors, it is also important to distinguish specific from nonspecific actions of OT. Recently, we have done the following experiment (Wang and Hatton, unpublished observations) to identify the receptors mediating OT functions in SON neurons. In the presence of an OT receptor antagonist, [β-Mercapto-β,β-cyclopentamethylene-propionyl]¹, O-Me-Tyr², Orn⁸]-oxytocin, the excitatory effects of on 6/6 OT neurons were blocked. However, a selective V₁ receptor antagonist, [deamino-Pen¹, O-Me-Tyr², Arg⁸]-vasopressin did not significantly influence

the excitatory effects of OT on any of the six OT neurons. Moreover, we have also tested the effect of a specific OT receptor antagonist (kindly provided by Dr. M. Manning) in one OT neuron, which also blocked OT effects. These results indicate that the excitatory effect of OT is achieved via activation of OT receptors.

Spike frequency reduction. Faced with progressively increasing OT concentrations and the resulting maintained membrane depolarization, SFR characterized the tonic firing activity of OT neurons. This is consistent with the actions of many other excitatory neuroactive substances (17). Distinct from spike frequency adaptation (17), the two major components leading to SFR in OT neurons, the firing rate decrease and E_m depolarization, are not regulated simultaneously. OT-evoked SFR could occur on the basis of dramatic depolarization (most active neurons) or subtle E_m changes (some less active neurons).

SFR was related to the resting E_m and the basal firing activity in most cases. In response to OT administration, E_m depolarized progressively, accompanied by firing rate increases. Firing rates, however, declined while changes in other parameters kept in the same direction as OT concentration rose to a certain level, suggesting that two separate mechanisms were at work, one primarily affecting firing activity, the other influencing spike features and E_m.

That OT receptors undergo rapid (within seconds to minutes) homologous desensitization after persistent agonist stimulation (9) may account for the SFR. Moreover, SFR may be related to activation of PKC and its negative feedback on the activation of phospholipase C. This may be explained by an activation of PKC that causes inhibition of subsequent responses of intracellular Ca^{2+} to activation of the diacylglycerol-PKC pathway (26). Other possible reasons are Ca^{2+} entry, which is important in E_m depolarization, and modulation of K^+ currents (1, 11) could cause decreased firing rate.

Local neural circuits and OT effects. In considering the role of neural circuits, glutamatergic inputs are undoubtedly involved. Besides reducing the depolarizing effect of OT, blocking ionotropic glutamate receptors also blocked excitatory responses of OT neurons to low doses of OT, and delayed or removed SFR. This is in agreement with *in vivo* experiments investigating the functions of glutamate (23). On the other hand, inhibiting the effects of inhibitory neurotransmitters also modulated OT actions, that is, bicuculline increased OT excitatory effect. The balance of glutamatergic and GABAergic actions is an important factor. Although OT reduced EPSCs (14), it also decreased IPSCs (3). Because GABAergic inhibitory inputs were dominant, the inhibition of presynaptic currents by OT shifted the excitatory-inhibitory balance, permitting a dominant excitatory innervation by glutamatergic neurons. Thus, through activating cationic currents (AMPA receptors) or Ca^{2+} currents (NMDA receptors), glutamate could cause depolarization and excitation.

Physiological implications of OT autoregulation. Under *in vivo* conditions, the basal firing rate of OT neurons in lactating rats is low, and OT fails to facilitate OT neuronal activity in the absence of suckling stimulation (18). Possibly, basal firing activity of OT neurons may have already adapted to basal OT levels in that situation, as the higher concentrations of exogenous OT produced only permissive effects for the actions of other neuroactive substances during suckling stimulation (5, 6). That OT actions were weakened by blocking ionotropic glutamate receptors and were facilitated during fast GABAergic blockade suggests the OT actions are under intense modulation of the neurochemical environment, and suckling may allow reestablishment of time- and concentration-dependent actions.

Upon nursing, suckling-associated neural pathways (e.g., noradrenergic, glutamatergic, and OTergeric) are mobilized (24). Such inputs result in somatodendritic release of OT without causing significant changes in basal firing rates (27, 16) or adjusting the activity of OT neurons at moderate firing rates (2) by SFR. In our results for prolonged application of 1 pM OT, one highly active neuron did not increase its firing frequency but rather decreased its firing rate in response to OT application, supporting a dual modulatory role of OT in OT neuronal activity. Increased OT release inhibits GABAergic input, while sensitizing responses to glutamate and noradrenaline, lead to the initial activation of OT neurons. This activation further increases local OT concentrations (8). OT will activate extracellular peptidases (14), accelerating OT decomposition. Astroglia in the SON will also help in the removal of excessive OT. In addition, nitric oxide and adenosine together with GABA will suppress the firing activity and aid in repolarization of E_m , resensitizing OT neurons to excitatory inputs. As a result, the firing rate increases gradually, and a new cycle of oscillations in the local neural circuits can begin.

In conclusion, progressively increasing OT concentration causes excitation and subsequent SFR, accompanied by a series of changes in other membrane electrical features. Glutamatergic, but not GABAergic, inputs are influential in the firing rate changes, whereas other membrane electrical changes are modulated mainly through postsynaptic processes. These effects may well reflect the actions of OT during suckling stimulation and represent a common working model of neuropeptides on their secretory neurons.

ACKNOWLEDGMENTS

We thank Dr. Harold Gainer at the National Institutes of Health for providing PS38 antibody and Dr. Maurice Manning for the specific OT antagonist.

GRANTS

Research was supported by National Institutes of Health Grant NS-009140.

REFERENCES

1. Bourque CW and Renaud LP. Calcium-dependent action potentials in rat supraoptic neurosecretory neurones recorded *in vitro*. *J Physiol* 363: 419–428, 1985.
2. Brown D and Moos F. Onset of bursting in oxytocin cells in suckled rats. *J Physiol* 503: 625–634, 1997.
3. Brussaard AB, Kits KS, and de Vlieger TA. Postsynaptic mechanism of depression of GABAergic synapses by oxytocin in the supraoptic nucleus of immature rat. *J Physiol* 497: 495–507, 1996.
4. Carter SJ, Richardson CM, and Wakerley JB. Excitatory effects of oxytocin and cholecystokinin on oxytocin neurons: differences between virgin, pregnant and lactating rats. *Neurosci Lett* 351: 13–16, 2003.
5. Cosgrave AS, Richardson CM, and Wakerley JB. Permissive effect of centrally administered oxytocin on the excitatory response of oxytocin neurones to ventral tegmental stimulation in the suckled lactating rat. *J Neuroendocrinol* 12: 843–852, 2000.
6. Cosgrave AS and Wakerley JB. Central oxytocin potentiates excitatory responses of oxytocin neurones to stimulation of the dorsomedial hypothalamic nucleus in the suckled rat. *Neurosci Lett* 317: 47–49, 2002.
7. Crowley WR and Armstrong WE. Neurochemical regulation of oxytocin secretion in lactation. *Endocr Rev* 13: 33–65, 1992.
8. de Kock CP, Wierda KD, Bosman LW, Min R, Koksma JJ, Mansvelder HD, Verhage M, and Brussaard AB. Somatodendritic secretion in oxytocin neurons is upregulated during the female reproductive cycle. *J Neurosci* 23: 2726–2734, 2003.
9. Evans JJ, Forrest-Owen W, and McArdle CA. Oxytocin receptor-mediated activation of phosphoinositidase C and elevation of cytosolic calcium in the gonadotrope-derived alphaT3–1 cell line. *Endocrinology* 138: 2049–2055, 1997.
10. Freund-Mercier MJ and Richard P. Electrophysiological evidence for facilitatory control of oxytocin neurones by oxytocin during suckling in the rat. *J Physiol* 352: 447–466, 1984.
11. Hatton GI and Li Z. Mechanisms of neuroendocrine cell excitability. *Adv Exp Med Biol* 449: 79–95, 1998.
12. Hatton GI and Yang QZ. Synaptic potentials mediated by alpha 7 nicotinic acetylcholine receptors in supraoptic nucleus. *J Neurosci* 22: 29–37, 2002.
13. Klein U and Fahrenholz F. Reconstitution of the myometrial oxytocin receptor into proteoliposomes. Dependence of oxytocin binding on cholesterol. *Eur J Biochem* 220: 559–567, 1994.
14. Kombian SB, Mougnot D, and Pittman QJ. Dendritically released peptides act as retrograde modulators of afferent excitation in the supraoptic nucleus *in vitro*. *Neuron* 19: 903–912, 1997.
15. Kow LM, Johnson AE, Ogawa S, and Pfaff DW. Electrophysiological actions of oxytocin on hypothalamic neurons *in vitro*: neuropharmacological characterization and effects of ovarian steroids. *Neuroendocrinology* 54: 526–535, 1991.
16. Ludwig M, Sabatier N, Bull PM, Landgraf R, Dayanithi G, and Leng G. Intracellular calcium stores regulate activity-dependent neuropeptide release from dendrites. *Nature* 418: 85–89, 2002.
17. McCormick DA. Membrane properties and neurotransmitter actions. In: *The Synaptic Organization of the Brain* (5th ed.), edited by Shepherd GM. New York: Oxford University Press, 2004, p. 39–77.

18. Meyer C, Freund-Mercier MJ, and Richard P. Facilitatory effect of oxytocin on oxytocin cell background activity in the rat is suckling-dependent. *Neurosci Lett* 75: 80–84, 1987.
19. Moos F, Freund-Mercier MJ, Guerne Y, Guerne JM, Stoeckel ME, and Richard P. Release of oxytocin and vasopressin by magnocellular nuclei in vitro: specific facilitatory effect of oxytocin on its own release. *J Endocrinol* 102: 63–72, 1984.
20. Moos F, Poulain DA, Rodriguez F, Guerne Y, Vincent JD, and Richard P. Release of oxytocin within the supraoptic nucleus during the milk ejection reflex in rats. *Exp Brain Res* 76: 593–602, 1989.
21. Neumann I, Koehler E, Landgraf R, and Summy-Long J. An oxytocin receptor antagonist infused into the supraoptic nucleus attenuates intranuclear and peripheral release of oxytocin during suckling in conscious rats. *Endocrinology* 134: 141–148, 1994.
22. Neumann I, Russell JA, and Landgraf R. Oxytocin and vasopressin release within the supraoptic and paraventricular nuclei of pregnant, parturient and lactating rats: a microdialysis study. *Neuroscience* 53: 65–75, 1993.
23. Parker SL and Crowley WR. Central stimulation of oxytocin release in the lactating rat by N-methyl-D-aspartate: requirement for coactivation through non-NMDA glutamate receptors or the glycine coagonist site. *Neuroendocrinology* 62: 467–478, 1995.
24. Parker SL and Crowley WR. Stimulation of oxytocin release in the lactating rat by a central interaction of alpha 1-adrenergic and alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic acid-sensitive excitatory amino acid mechanisms. *Endocrinology* 133: 2855–2860, 1993.
25. Stern JE and Armstrong WE. Sustained outward rectification of oxytocinergic neurones in the rat supraoptic nucleus: ionic dependence and pharmacology. *J Physiol* 500: 497–508, 1997.
26. Tornquist K and Tashjian AH Jr. 12-O-tetradecanoyl-phorbol-13-acetate decreases influx of extracellular Ca^{2+} induced by depolarization in GH4C1 cells: effects of pretreatment with 1,25-dihydroxycholecalciferol. *Endocrinology* 126: 2068–2078, 1990.
27. Wakerley JB, Clarke G, and Summerlee AJ. Milk ejection and its control. In: *The Physiology of Reproduction* (2nd ed.), edited by Knobil E and Neill JD. New York: Raven Press, 1994, p. 1131–1177.
28. Yamashita H, Okuya S, Inenaga K, Kasai M, Uesugi S, Kannan H, and Kaneko T. Oxytocin predominantly excites putative oxytocin neurons in the rat supraoptic nucleus in vitro. *Brain Res* 416: 364–368, 1987.

