# Prenatal blockade of estradiol synthesis impairs respiratory and metabolic responses to hypoxia in newborn and adult rats

V. D. Doan, S. Gagnon and V. Joseph

*Am J Physiol Regul Integr Comp Physiol* 287:R612-R618, 2004. First published 13 May 2004; doi: 10.1152/ajpregu.00627.2003

# You might find this additional info useful...

- This article cites 44 articles, 11 of which you can access for free at: http://ajpregu.physiology.org/content/287/3/R612.full#ref-list-1
- This article has been cited by 3 other HighWire-hosted articles: http://ajpregu.physiology.org/content/287/3/R612#cited-by
- Updated information and services including high resolution figures, can be found at: http://ajpregu.physiology.org/content/287/3/R612.full

Additional material and information about American Journal of Physiology - Regulatory, Integrative and Comparative Physiology can be found at: http://www.the-aps.org/publications/ajpregu

This information is current as of July 28, 2016.

*American Journal of Physiology - Regulatory, Integrative and Comparative Physiology* publishes original investigations that illuminate normal or abnormal regulation and integration of physiological mechanisms at all levels of biological organization, ranging from molecules to humans, including clinical investigations. It is published 12 times a year (monthly) by the American Physiological Society, 9650 Rockville Pike, Bethesda MD 20814-3991. Copyright © 2004 the American Physiological Society. ISSN: 0363-6119, ESSN: 1522-1490. Visit our website at http://www.the-aps.org/.

# Prenatal blockade of estradiol synthesis impairs respiratory and metabolic responses to hypoxia in newborn and adult rats

# V. D. Doan, S. Gagnon, and V. Joseph

Department of Pediatrics, Laval University, Centre de Recherche (D0–711), Hôpital St-François d'Assise, Quebec, Quebec G1L 3L5, Canada

Submitted 27 October 2003; accepted in final form 7 May 2004

Doan, V. D., S. Gagnon, and V. Joseph. Prenatal blockade of estradiol synthesis impairs respiratory and metabolic responses to hypoxia in newborn and adult rats. Am J Physiol Regul Integr Comp Physiol 287: R612-R618, 2004. First published May 13, 2004; 10.1152/ajpregu.00627.2003.-We tested the hypothesis that estradiol modifies respiratory control in pregnant rats and participates in the development of respiratory chemoreflexes in fetuses. Pregnant rats (n = 12) received daily subcutaneous injections of vehicle (Veh, n =6) or 4-androsten-4-ol-3,17-dione acetate (ATD; inhibitor of estradiol synthesis; n = 6; 5 mg/day in vehicle) from gestational day 16 (G16) to delivery. Baseline ventilation (whole body plethysmography) and metabolic rate [oxygen consumption (Vo<sub>2</sub>)] were determined at G14 and G20, in pups [on postnatal day 3 (P3) and P20] and in adult rats (on P70) born to Veh- or ATD-treated mothers. Hypoxic chemoreflex was assessed in P3 rats by acute exposure to 60% O2 and in P20 or P70 rats by moderate hypoxia (12% O2, 30 min). ATD treatment reduced circulating estradiol in pregnant dams at G20 without producing changes in the circulating level of estradiol precursors (testosterone and androstenedione). ATD-treated dams showed impaired respiratory adjustment to late gestation. Pups born to ATD mothers had higher resting Vo2 (+23% at P3, +21% at P20), respiratory frequency (+15% at P3, +12% at P20), and minute ventilation (+11% at P3, +18% at P20) than pups from Veh mothers. Respiratory decrease during acute hyperoxic exposure at P3 was -9.7% in Veh (P < 0.05 vs. room air) and only -2.6% (P = not significant) in ATD pups. In P20 ATD rats, hypoxic ventilatory response was attenuated compared with Veh. In P20 and P70 rats, the drop of Vo2 in hypoxia (-31% in P70, P < 0.0001) was not observed in ATD rats. We conclude that estradiol secreted during late gestation is necessary for respiratory adjustment to pregnancy and is required for adequate development of respiratory and metabolic control in the offspring.

hypoxic ventilatory response; prenatal estradiol synthesis; aromatase inhibitor; development

OVARIAN STEROIDS, including progesterone and estradiol, have long been recognized as specific factors able to modulate normoxic breathing and hypoxic chemoreflexes by acting both on the central nervous system (4-6) and on the dopaminergic signaling pathway in the peripheral chemoreceptors (21, 22, 40). These animal models are clinically relevant because the combined action of estradiol and progesterone on respiratory control is thought to reduce the occurrence of respiratory disorders such as sleep-disordered breathing (10, 36, 38, 47) and high-altitude deacclimatization (22, 28) in women.

During pregnancy, the circulating levels of estradiol and progesterone in maternal circulation increase by  $\sim$ 100-fold (33) as a result of the endocrine activity of the placenta (in

primate; Ref. 24) or ovaries (in rodents; Ref. 1), influencing both maternal and fetal physiology (16, 34). Additionally, estradiol synthesized directly in specific fetal brain neurons (i.e., dopaminergic nigrostriatal neurons) is a potent autocrine regulator of neural growth and survival, involving both classical (nuclear) and nonclassical (membrane bound) estradiol receptors (26).

Former studies emphasized the relationship between pregnancy-related hormonal secretion, increased minute ventilation (VE), and enhanced respiratory chemoreflexes observed during late gestation in humans (33). However, the role of estradiol exposure on the respiratory control of the fetus and newborn remains unknown. This particular aspect may be of clinical importance as preterm and extreme preterm births are characterized both by respiratory disorders (19, 27) and chronic deficit of estradiol normally provided by the endocrine placenta (41-43). Thus we hypothesized that gestational blockade of estradiol synthesis in rats would impair maternal respiratory adjustment to pregnancy and potentially disrupt the adequate development of respiratory control in newborn rats. We used a highly specific pharmacological blocker of estradiol synthesis [4-androsten-4-ol-3,17 dione acetate (ATD), a potent inhibitor of P-450 aromatase enzyme, the key enzyme for estradiol synthesis] (37) in pregnant female rats to effectively reduce the circulating level of estradiol during the last days of gestations. Our results are consistent with a yet unrecognized role of endogenous estradiol for adequate development of ventilatory control in newborn rats.

#### MATERIALS AND METHODS

Animal treatment. All experiments have been approved by the local committee (Laval University) of animal care and use. On the day of estrus (determined on the basis of daily vaginal smears), female Sprague-Dawley rats (n = 12; purchased from Charles River Canada) were put overnight with mating males. Pregnancy was confirmed after observation of spermatozoids in the vaginal smear the next day [considered as day 0 of gestation; G0] and weight gain during the subsequent days. Pregnant female rats were housed separately and left undisturbed until G14, when they were weighed and baseline ventilation and oxygen consumption (Vo<sub>2</sub>) were assessed (see below). Forty-eight hours later (G16), pregnant dams were divided into two groups receiving daily subcutaneous injections (500 µl) of vehicle (Veh; n = 6, 10% ethanol in propylene glycol) or the inhibitor of estradiol synthesis, ATD (n = 6; 5 mg/day in vehicle, Sigma-Aldrich) until delivery. At G20, females were weighed, and baseline respiratory and metabolic measurements were repeated.

Address for reprint requests and other correspondence: V. Joseph, Centre de Recherche CHUQ, Hôpital St. François d'Assise, Local D0–711, 10 Rue de l'Espinay, G1L 3L5 Quebec, Quebec, Canada (E-mail: joseph.vincent@ crsfa.ulaval.ca).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "*advertisement*" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

# PRENATAL ESTRADIOL AND DEVELOPMENT OF RESPIRATORY CONTROL

Table 1. Circulating levels of estradiol, testosterone, and androstenedione assessed by gas chromatography at gestational day 20 in Veh- and ATD-treated mothers

	Veh	ATD
Estradiol, pg/ml Testosterone, ng/ml Androstenedione, ng/ml	$28.1 \pm 2.6$ $0.40 \pm 0.08$ $0.40 \pm 0.08$	Non detectable (<5 pg/ml) 0.34±0.08 0.56±0.06

Values are means  $\pm$  SE. Since estradiol values in the 4-androsten-4-ol-3,17dione acetate (ATD)-treated group were below the detection threshold (5 pg/ml), statistical analysis was not performed for estradiol. Veh, vehicle.

After respiratory recordings, females were briefly anesthetized under halothane, and a sample of blood (500  $\mu$ l) was drawn from the jugular vein. To assess the efficiency of ATD treatment, hormonal assays were done by gas chromatography-mass spectrometry using negative chemical ionization to determine circulating levels of estradiol and its precursors (testosterone, androstenedione; Laboratory of Clinical Endocrinology, CHUL, University Laval, Quebec, Canada). Steroids were extracted from serum by liquid-liquid and solid-phase extraction, and derivatization reactions were performed to improve chromatographic and detection response as previously described (7).

At birth, male pups from all litters were kept for further analysis; females pups were used to normalize the size of each litter to 12 pups to avoid litter size effects on postnatal growth. All the animals (males and females) were weighed between 24 and 36 h after birth, and pups were raised with their natural mothers until weaning.

Respiratory recordings in pregnant dams and offspring. Respiratory recordings were performed by whole body plethysmography (Emka Technologies, Paris, France) using a method derived from the pressure plethysmograph described by Bartlett and Tenney (3), with chambers of different size for adults and pups. The recording chamber included a built-in reference chamber and was permanently flushed with fresh air (or the desired gas mixture; see below). The Emka chamber allowed measurements of flow related to respiratory gas exchanges in the chamber. After adequate calibration of the chamber, all measurements were corrected with the standard equation (3) to allow determination of tidal volume after integration of the flow signal by the Emka software. Temperature and relative humidity inside the chamber were constantly monitored during recordings, and barometric pressure was read from a barometer. For 3-day-old rats, body temperature was considered to be stable at 35°C with a mean ambient temperature of  $30.3 \pm 0.08$  °C inside the recording chamber.

The inlet air was circulated at a regular flow of 400 ml/min [postnatal *day 3* (P3)], 600 ml/min (weaning), or 1,500 ml/min (adults) through a dedicated oxygen sensor (s103 O<sub>2</sub> analyzer, Qubit-systems, Kingston, ON, Canada) and a subsampling pump and drying column allowed the outlet flow to be monitored by a second oxygen sensor for determination of  $\dot{V}O_2$  [( $\%O_{2 in} - \%O_{2 out}$ ) × flow]. The oxygen sensors were calibrated daily before each experiments. The outlet-subsampling pump was switched on only for metabolic measurements, which lasted 2–3 min and were initiated after 10–15 min of stable ventilatory recordings and at the end of the hypoxic exposure. For each animal, at the end of the hypoxic exposure of the animal had been removed from the chamber to ensure adequate baseline readings in hypoxia. All signals (ventilation and O<sub>2</sub>%) were recorded and stored on a computer for later analysis.

Basal ventilation and respiratory responses to hyperoxia in newborn pups (Dejours test). Respiratory measurements were done in 3-day-old (P3) male pups born to Veh (n = 15)- or ATD-treated (n =18) mothers. After a period of habituation in the recording chamber (15–30 min) and at least 10 min of stable baseline recordings, the inlet flow was derived from a hyperoxic gas mixture, which allowed a rapid exposure to 60% O<sub>2</sub> in the chamber. Respiratory changes were analyzed during the first 20 s of hyperoxic exposure. This procedure, classically referred to as "Test de Dejours" (18, 23), is highly effective to address the sensitivity of peripheral chemoreceptors in vivo.

In weanling and adult rats, the same procedure for baseline recordings was repeated, and then the inlet flow was derived from a hypoxic gas mixture allowing rapid exposure to 12% O<sub>2</sub> for 30 min. The respiratory response to hypoxia was analyzed minute by minute during the first 10 min of exposure and as the average of the last 5 min of exposure. Rectal temperature was measured before the onset of normoxic recordings and at the end of the hypoxic exposure with a thermocouple thermometer. We used a total of 15 and 18 weanling rats from Veh- and ATD-treated dams, respectively, and 11 and 15 adults from Veh and ATD dams.

Statistical analysis. For simple measurements (1 factor), data were analyzed by one-way ANOVA. For longitudinal comparisons, data were analyzed by two-way ANOVA for repeated measures. If a global effect or significant interaction appeared between groups on the general ANOVA, data were analyzed for factorial measures by a post hoc Fisher's protected least significant difference test. The results are indicated in the corresponding tables and by symbols in the figures. All values are means  $\pm$  SE, and the level of significance was set at 0.05.

# RESULTS

Blockade of estradiol synthesis impairs respiratory adaptation during late gestation in female rats. The ATD treatment reduced circulating estradiol measured at G20 in pregnant dams to undetectable levels (<5 pg/ml) without producing significant increases in the circulating level of the substrates of the P-450 aromatase, i.e., testosterone and androstenedione (see Table 1). Between G14 and G20. Veh- and ATD-treated dams had similar increase in body weight (from  $363 \pm 11$  to 432  $\pm$  14 g in Veh-treated and 377  $\pm$  6 to 452  $\pm$  9 g in ATD-treated dams), but the normal increase of tidal volume that takes place during late gestation in Veh-treated mothers (from 1.85  $\pm$  0.09 to 2.35  $\pm$  0.07 ml at G14 and G20, respectively) was blocked in ATD-treated dams (1.96  $\pm$  0.10 ml at G14 and 1.91  $\pm$  0.13 ml at G20, Fig. 1). Nonetheless, G20 ATD-treated dams maintained  $\dot{V}_E$  (37.8  $\pm$  2.9 ml·min<sup>-1</sup>·100 g<sup>-1</sup>) at a level similar to Veh-treated mothers  $(41.0 \pm 0.5 \text{ ml} \cdot \text{min}^{-1} \cdot 100 \text{ g}^{-1})$  by slightly increasing respiratory frequency (from  $81.3 \pm 2.5$  at G14 to  $90.1 \pm 6.4$ breaths/min at G20), while at the same time Veh-treated dams slightly decreased their respiratory frequency (from  $82.7 \pm 3.8$ to 75.2  $\pm$  1.6 breaths/min).



Fig. 1. Evolution of respiratory parameters between gestational *days* 14 and 20 in dams treated with vehicle (Veh) or 4-androsten-4-ol-3,17-dione acetate (ATD). BW, body weight; V<sub>t</sub>, tidal volume; F<sub>r</sub>, respiratory frequency;  $\dot{V}_E$ , minute ventilation. Results are expressed as means  $\pm$  SE of %changes between gestational *day* 14 (G14) and G20 calculated for each dam. # P < 0.05, G20 vs. G14: \*P < 0.05, ATD vs. Veh.

PRENATAL ESTRADIOL AND DEVELOPMENT OF RESPIRATORY CONTROL



Fig. 2. Effects of prenatal ATD on baseline values for body weight (*A*), respiratory frequency (*B*), tidal volume (*C*), minute ventilation (*D*), oxygen consumption ( $\dot{V}o_2$ ; *E*), and  $\dot{V}E/\dot{V}o_2$  ratio in 3-, 20-, or 70-day-old rats. For each age, values from rats born to Veh dams are considered 100% (mean  $\pm$  SE). \**P* < 0.05, ATD vs. Veh.

There was a similar decrease of  $\dot{V}o_2$  between G14 and G20 in Veh (from 2.74 ± 0.16 to 2.14 ± 0.26 ml·min<sup>-1</sup>·100 g<sup>-1</sup>, P < 0.05) and ATD (from 2.60 ± 0.25 to 2.00 ± 0.13 ml·min<sup>-1</sup>·100 g<sup>-1</sup>, P < 0.05), leading to a similar increase of  $\dot{V}E/\dot{V}o_2$  between G14 and G20 for the two groups of dams (from 15.8 ± 1.3 to 20.1 ± 2.3 for Veh and from 15.5 ± 1.2 to 18.9 ± 0.9 for ATD dams). At birth, males and females born to ATD dams had reduced body weight (6.49 ± 0.17 vs. 6.96 ± 0.11 g for males, 6.29 ± 0.08 vs. 6.81 ± 0.11 g for females, P < 0.0001, ATD vs. Veh for both gender).

Gestational blockade of estradiol synthesis produces longlasting alterations of respiratory and metabolic control in normoxic control conditions. Metabolic and respiratory results under normoxic conditions in rats born to ATD- and Vehtreated mothers are compiled in Fig. 2, A-F. Three-day-old male pups born to mothers treated with ATD had a lower body weight (8.53  $\pm$  0.22 g; n = 18) compared with pups from Veh-treated dams (9.50  $\pm$  0.23 g; n = 15; P = 0.006, Fig. 2A) that was no longer apparent in weanling (ATD 50.1  $\pm$  1.4 g vs. Veh 52.2  $\pm$  1.4 g) and adult rats (ATD 476  $\pm$  10 g vs. Veh  $468 \pm 12$  g; Fig. 2A). Compared with rats born to Veh-treated mothers, rats from ATD mothers presented higher respiratory frequency at 3 and 20 days of age  $(171 \pm 5 \text{ vs. } 149 \pm 4 \text{ and}$  $148 \pm 3$  vs.  $132 \pm 4$  breaths/min, P = 0.002 for both ages, Fig. 2B), higher  $\dot{V}_E$  in 3-day-old rats (157  $\pm$  6 vs. 141  $\pm$  6  $\text{ml}\cdot\text{min}^{-1}\cdot100 \text{ g}^{-1}$ , P = 0.05, Fig. 2D) and a tendency toward higher level in 20-day-old rats also (126  $\pm$  4 vs. 100  $\pm$  7 ml·min<sup>-1</sup>·100 g<sup>-1</sup>, P = 0.07), and increased  $\dot{V}o_2$  at 3 and 20 days of age (7.38  $\pm$  0.35 vs. 6.00  $\pm$  0.43 and 4.43  $\pm$  0.24 vs.  $3.65 \pm 0.24 \text{ ml} \cdot \text{min}^{-1} \cdot 100 \text{ g}^{-1}$ ; P = 0.02 and 0.04, respectively, Fig. 2*E*). The corresponding  $\dot{V}_E/\dot{V}_{O_2}$  was not altered at any age after ATD treatment (Fig. 2F). Adult rats born to ATD-treated dams had lower tidal volume under normoxia than rats from Veh-treated mothers (0.32  $\pm$  0.01 vs. 0.39  $\pm$ 0.02 ml/100 g, P = 0.008, Fig. 2C), but they had similar  $\dot{V}_E$   $(35.1 \pm 2.5 \text{ vs. } 39.4 \pm 1.1 \text{ ml} \cdot \text{min}^{-1} \cdot 100 \text{ g}^{-1}; P = 0.2, \text{ Fig.}$ 2D),  $\dot{V}o_2$  (2.08  $\pm$  0.10 vs. 2.33  $\pm$  0.10 ml $\cdot$ min $^{-1} \cdot 100 \text{ g}^{-1};$ P = 0.1, Fig. 2E), and  $\dot{V}E/\dot{V}o_2$  ratio (18.3  $\pm$  1.6 vs. 17.7  $\pm$  1.6; P = 0.8, Fig. 2F).

Gestational blockade of estradiol synthesis produces longlasting alterations of hypoxic ventilatory and metabolic responses. When exposed to a brief hyperoxic period (Dejours test), 3-day-old pups born to Veh-treated mothers decreased  $\dot{V}_E$ (from 142 ± 6 to 127 ± 3 ml·min<sup>-1</sup>·100 g<sup>-1</sup>, P < 0.02) and respiratory frequency (from 148 ± 4 to 117 ± 4 breaths/min, P < 0.0001), while ATD pups had a reduced response ( $\dot{V}_E$ from 159 ± 6 to 154 ± 6 ml·min<sup>-1</sup>·100 g<sup>-1</sup>, P = not significant; respiratory frequency from 172 ± 5 to 148 ± 4, P < 0.0001) compared with Veh (see Fig. 3).

In weanling rats, the hypoxic ventilatory response was blunted by prenatal blockade of estradiol synthesis (Figs. 4 and 5); this pattern was not apparent in adults (Figs 4 and 5). At the end of the hypoxic exposure (12% O<sub>2</sub>; 30 min), weanling rats



Fig. 3. Percent changes of respiratory variables ( $\dot{V}E$  and  $F_r$ ) during the first 20 s of hyperoxic exposure in newborn rats born to Veh- or ATD-treated mothers. # P < 0.05, hyperoxia vs. normoxia; \*P < 0.05, ATD vs. Veh.



Fig. 4. Normalized minute-by-minute hypoxic ventilatory response in weanling [postnatal *day 20* (P20), *left*] and adult rats (P70, *right*) born to Veh- or ATD-treated mothers during 10 min of hypoxic exposure (12% O<sub>2</sub>). Ve (ml·min<sup>-1</sup>·100 g<sup>-1</sup>), V<sub>t</sub> (ml/100 g), and F<sub>r</sub> (breaths/ min) are expressed as % of baseline values. All values are means  $\pm$  SE. \**P* < 0.05, ATD vs. Veh.



groups) or in adults  $(38.1 \pm 0.1^{\circ}\text{C} \text{ in both groups})$ , and the magnitude of rectal temperature drop after 30 min of hypoxic exposure was similar in ATD or Veh rats (~1°C).

#### DISCUSSION

The results of this study are consistent with the admitted hypothesis that the increase of circulating estradiol during late gestation participates in the respiratory adjustments observed during late pregnancy (13, 33) and demonstrate that fetal exposure to estradiol is required for normal development of respiratory and metabolic control. In newborn, weanling, and adult rats born to ATD-treated mothers, the alterations of respiratory and metabolic control included *1*) increased resting  $\dot{V}o_2$ , 2) increased respiratory frequency and  $\dot{V}E$ , 3) decreased



Fig. 5. Percent changes of metabolic and respiratory parameters between normoxia and hypoxia exposure (12% O<sub>2</sub>; 30 min) in weanling (P20; *left*) and adult (P70; *right*) rats born to Veh- or ATD-treated dams. Vo<sub>2</sub> (ml·min<sup>-1</sup>·100 g<sup>-1</sup>),  $\dot{V}_{E}$  (ml·min<sup>-1</sup>·100 g<sup>-1</sup>), and  $\dot{V}_{E}/\dot{V}_{O_2}$  ratio. H P < 0.05, hypoxia vs. normoxia; \*P < 0.05, ATD vs. Veh.

AJP-Regul Integr Comp Physiol • VOL 287 • SEPTEMBER 2004 • www.ajpregu.org

sensitivity of the peripheral chemoreceptors (assessed by Dejours test in newborn rats), and 4) altered metabolic response to hypoxia in weanling and adult rats. These effects reflect the potential diversity of action of steroid hormones such as estradiol on fetal respiratory and metabolic control development.

Methodological considerations. We used a derivate (ATD) of androstenedione that has been specifically designed as a potent and selective inhibitor of P-450 aromatase enzyme (the key enzyme for estradiol synthesis) and that irreversibly binds to the active site of the enzyme as a competitor to endogenous substrate (37). The P-450 aromatase enzyme is a cytochrome protein converting testosterone to estradiol and androstenedione to estrone and is expressed at high level during late gestation in the placenta of primate (24), in the ovaries of rodent (1), and in some discrete brain areas of rat fetuses, where estradiol acts as a specific developmental factor (26, 44). ATD is highly effective in reducing the circulating levels of estradiol (12, 37) as well as immunostaining of brain aromatase (17) and aromatase mRNA concentration in hypothalamic preoptic area in quail (2). Because ATD binds irreversibly to the aromatase, recovery of complete estradiol synthesis is not completed until new aromatase protein is synthesized (37). In humans, after a single injection of an aromatase inhibitor showing similar properties (4-hydroxyandrostene-3,17-dione, 500 mg), the level of circulating estradiol is decreased by 50% for at least 7 consecutive days (12). Thus, if ATD crosses the placenta, it is highly probable that its effect will be maintained for several postnatal days in rat pups. Our own results show that in pregnant dams, ATD can be used to reduce the circulating levels of estradiol without producing an increase in the circulating levels of androstenedione and testosterone, the direct precursors for estradiol synthesis. On the other hand, this pharmacological model does not allow us to differentiate between the influences of maternal estradiol furnished to the fetus and in situ estradiol synthesis in fetal neurons (see below for further details). Nonetheless, this model is particularly interesting because it describes for the first time the role of endogenously secreted estradiol for the development of respiratory reflexes. Whether this is related to maternofetal estradiol transfer or depends on local (i.e., fetal) estradiol synthesis will require further experiments.

Effects of ATD in pregnant dams. Pregnant dams treated with ATD demonstrated consistent signs of impaired respiratory adjustments, including lower tidal volume compared with Veh-treated mothers, which is in good agreement with previous studies showing that ovarian steroids stimulate breathing almost exclusively by increasing tidal volume (22, 25, 35). This effect is mediated through central areas, including hypothalamic nuclei (6), and by a reduction of the dopaminergic inhibitory drive at the level of the peripheral chemoreceptors (22). The central facilitation of breathing by ovarian steroids requires the induction of progesterone receptors by estradiol in central areas (4-6). The present results clearly emphasize the critical importance of the stimulatory effects of estradiol for respiratory adjustments during late gestation in rats. Contrastingly, pups born to ATD-treated dams had specific alterations, thus revealing that estradiol synthesis is an important factor for adequate respiratory development.

*Effects of ATD in pups.* Three-day-old male pups born to treated mothers had increased metabolic rate and elevated

respiratory frequency and VE compared with pups from Vehtreated dams. The similar VE/VO2 ratio in treated pups compared with controls indicates that the direct drive of basal metabolic rate on breathing was not affected by prenatal hormonal blockade. Therefore, the increased VE likely results from the higher metabolic drive observed in treated pups. On the other hand, the decreased respiratory responses to acute hyperoxia (in newborn) and hypoxia (in weanling rats) in ATD pups strongly suggests that the responsiveness of the peripheral chemoreceptors to hypoxia has been altered. Nonetheless, and because a metabolic effect on respiratory integration cannot be excluded at that time, we will require more direct evidence to sustain (or not) this hypothesis. The metabolic and ventilatory alterations under normoxic or hypoxic conditions persisted until weaning (respiratory) and adulthood (metabolic), suggesting that these were not short-lasting effects of estradiol deficit, but rather long-term (or permanent) alterations of the respiratory and metabolic regulatory systems.

Developmental regulation of respiratory and metabolic control by estradiol. Direct fetal exposure to estradiol can occur either as the result of placental transfer from maternal circulation to the fetus or from local synthesis of estradiol in fetal brain. P-450 aromatase mRNA is expressed at high levels in the ovaries of pregnant female rats starting from G8 (1). In the fetal brain, immunohistochemical staining of P-450 aromatase protein appears at G13 in the hypothalamic preoptic area and is sustained at high level in hypothalamic nuclei from G16 to P7 in rats (44, 45). During this developmental window, testosterone (the endogenous precursor for estradiol synthesis by P-450 aromatase) can be provided to the fetus by the activity of testes in males (16) or as the result of local neurosteroid synthesis (48). Recent clues have accumulated showing that prenatal estradiol plays a crucial role in fetal development: estradiol receptors are expressed in numerous tissues in middle-gestation human fetuses (11), estradiol accelerates fetal lung maturation by stimulation of surfactant phospholipids synthesis (34), and, in fetal mice lacking the  $\beta$ -isoform of estrogen receptor, survival and migration of cortical neurons are both severely impaired during late gestation (46). Estradiol receptors (divided into nuclear and membrane bound) are located in a variety of neural populations, including hypothalamic and brain stem monoaminergic nuclei (see Refs. 8, 31, 32), which participate in ventilatory control under normoxic or hypoxic exposure (9, 14, 15, 20, 29) and may also contribute to the respiratory changes observed after prenatal blockade of estradiol synthesis.

Nonetheless, and because estradiol exerts a wide variety of action on different targets during development, it is difficult for the moment to propose a mechanistic framework to explain our results. The normal pattern of ventilation is generated in the pre-Bötzinger complex of the lower brain stem (39) and is modulated by a variety of neural and visceral afferent messages emerging from central and peripheral chemoreceptors, pulmonary receptors (30), and suprapontic nuclei located for example within the hypothalamus (15). While the results of the acute hyperoxic exposure in newborn rats suggest an impairment of peripheral chemoreceptor activity in pups born to ATD-treated dams, we presently cannot further speculate on the origins of respiratory and metabolic changes observed in rats born to ATD-treated dams.

Our present findings may have valuable clinical implications. Recent studies emphasized the potential therapeutic benefits of replacement of circulating estradiol and progesterone at normal levels (such as observed in utero or immediately after delivery) in extreme preterm infants that are otherwise chronically deprived from the hormonal placental milieu (41– 43). As discussed above, at that time our model cannot be used to discriminate between the maternofetal transfer and local fetal synthesis of estradiol on respiratory control system but describes for the first time the important developmental role of adequate prenatal estradiol synthesis for respiratory and metabolic control development.

#### ACKNOWLEDGMENTS

We acknowledge Dr. A. Bélanger and R. Bérubé (CHUL Research Center, Laval University, Quebec) for hormonal assays.

#### GRANTS

This study has been founded by The Foundation for Sick Children's Hospital (Grant XG 04–032), Fond de la Recherche en Santé du Québec, Association Pulmonaire du Québec, Fondation de la Recherche des Maladies Infantiles, Fondation du Centre Hôspitalo-Universitaire de Québec, and Chaire de Recherche Jeanne et Jean-Louis Lévesque.

#### REFERENCES

- Akinola LA, Poutanen M, Vihko R, and Vihko P. Expression of 17β-hydroxysteroid dehydrogenase type 1 and type 2, P450 aromatase, and 20α-hydroxysteroid dehydrogenase enzymes in immature, mature, and pregnant rats. *Endocrinology* 138: 2886–2892, 1997.
- Balthazart J. Steroid control and sexual differentiation of brain aromatase. J Steroid Biochem Mol Biol 61: 323–339, 1997.
- Bartlett D and Tenney SM. Control of breathing in experimental anemia. *Respir Physiol* 10: 384–395, 1970.
- Bayliss DA, Cidlowski JA, and Millhorn DE. The stimulation of respiration by progesterone in ovariectomized cat is mediated by an estrogen-dependent hypothalamic mechanism requiring gene expression. *Endocrinology* 126: 519–527, 1990.
- Bayliss DA and Millhorn DE. Central neural mechanisms of progesterone actions: application to the respiratory system. *J Appl Physiol* 73: 393–404, 1992.
- Bayliss DA, Millhorn DE, Gallman EA, and Cidlowski JA. Progesterone stimulates ventilation through a central nervous system steroid receptor-mediated mechanism in the cat. *Proc Natl Acad Sci USA* 84: 7788– 7792, 1987.
- Bérubé R, Malenfant J, Gauvin D, Blais E M, Gagnon R, Dumas MC, Racine J, Bourque J, and Bélanger A. Quantitation of androgenic and estrogenic steroids in human serum using gas chromatography and negative chemical ionization mass spectrometry. *Proc 48th ASMS Conf Mass Spectrometry Allied Topics*, Long Beach, California, 2000.
- Beyer C and Karolczak M. Estrogenic stimulation of neurite growth in midbrain dopaminergic neurons depends on cAMP/protein kinase A signalling. J Neurosci Res 59: 107–116, 2000.
- Bianchi AL, Denavit-Saubié M, and Champagnat J. Central control of breathing in mammals: neuronal circuitry, membrane properties and neurotransmitters. *Physiol Rev* 75: 1–45, 1995.
- Block A, Boysen P, Wynne J, and Hunt L. Sleep apnea, hypopnea and oxygen desaturation in normal subjects. A strong male predominance. *N Engl J Med* 300: 513–517, 1979.
- Brandenberger AW, Tee MK, Lee JY, Chao V, and Jaffe RB. Tissue distribution of estrogen receptors α (ER-α) and β (ER-β) mRNA in the midgestational human fetus. *J Clin Endocrinol Metab* 82: 3509–3512, 1997.
- Brodie AM, Wing LY, Goss P, Dowsett M, and Coombes RC. Aromatase inhibitors and the treatment of breast cancer. *J Steroid Biochem* 24: 91–97, 1986.
- 13. Brownell LG, West P, and Kryger MH. Breathing during sleep in normal pregnant women. *Am Rev Respir Dis* 133: 38–41, 1986.
- Dreshaj IA, Haxhiu MA, and Martin RJ. Role of the medullary raphe nuclei in the respiratory response to CO<sub>2</sub>. *Respir Physiol* 111: 15–23, 1998.

- Dreshaj IA, Haxhiu MA, Martin RJ, and Young JK. The basomedial hypothalamus modulates the ventilatory response to hypoxia in neonatal rats. *Pediatr Res* 53: 945–949, 2003.
- Fisher DA. Endocrinology of fetal development. In: Williams Textbook of Endocrinology (10th ed.), edited by Larsen PD, Kronenberg HM, Melmed S, and Polonsky KS. Philadelphia, PA: Saunders, 2003, p. 811–841.
- Foidart A, Tlemcani O, Harada N, Abe-Dohmae S, and Balthazart J. Pre- and post-translational regulation of aromatase by steroidal and nonsteroidal aromatase inhibitors. *Brain Res* 701: 267–278, 1995.
- Gautier H. Honoring Pierre Dejours: his contribution to the study of the role of arterial chemoreceptors in the regulation of breathing in humans. In: *Chemoreception: From Cellular Signaling to Functional Plasticity*, edited by Pequignot JM, Gonzales C, Nurse C, Prabhakar NR, and Dalmaz Y. New York: Kluwer Academics/Plenum, 2003, p. 1–7.
- Henderson-Smart DJ. The effect of gestational age on the incidence and duration of recurrent apnoea in newborn babies. *Aust Paediatr J* 17: 273–276, 1981.
- Joseph V and Pequignot JM. Neurochemical processes involved in acclimatization to long term hypoxia. In: Oxygen Sensing: Responses and Adaptation to Hypoxia, edited by Lahiri S, Semenza GL, and Prabhakar NR. New York: Dekker, 2003, p. 467–488.
- Joseph V, Soliz J, Pequignot J, Sempore B, Cottet-Emard JM, Dalmaz Y, Favier R, Spielvogel H, and Pequignot JM. Gender differentiation of the chemoreflex during growth at high altitude: functional and neurochemical studies. *Am J Physiol Regul Integr Comp Physiol* 278: R806–R816, 2000.
- 22. Joseph V, Soliz J, Soria R, Pequignot J, Favier R, Spielvogel H, and Pequignot JM. Dopaminergic metabolism in carotid bodies and high altitude acclimatization in female rats. *Am J Physiol Regul Integr Comp Physiol* 282: R765–R773, 2002.
- 23. Kline DD, Peng YJ, Manalo DJ, Semenza GL, and Prabhakar NR. Defective carotid body function and impaired ventilatory responses to chronic hypoxia in mice partially deficient for hypoxia-inducible factor 1α. Proc Natl Acad Sci USA 99: 821–826, 2002.
- 24. Kragie L. Aromatase in primate pregnancy: a review. *Endocr Res* 28: 121–128, 2002.
- 25. Kryger M, McCullough RE, Collins DD, Scoggin CH, Weil JV, and Grover RF. Treatment of excessive polycythemia of high altitude with respiratory stimulant drugs. *Am Rev Respir Dis* 117: 455–464, 1978.
- Kuppers E, Ivanova T, Karolczak M, and Beyer C. Estrogen: a multifunctional messenger to nigrostriatal dopaminergic neurons. *J Neurocytol* 29: 375–385, 2000.
- Lee D, Caces R, Kwiatkowski K, Cates D, and Rigatto H. A developmental study on types and frequency distribution of short apneas (3 to 15 seconds) in term and preterm infants. *Pediatr Res* 22: 344–349, 1987.
- León-Velarde F, Ramos MA, Hernández JA, De Idiáquez D, Muñoz LS, Gaffo A, Córdova S, Durand D, and Monge-CC. The role of menopause in the development of chronic mountain sickness. Am J Physiol Regul Integr Comp Physiol 272: R90–R94, 1997.
- Mack SO, Kc P, Wu M, Coleman BR, Tolentino-Silva FP, and Haxhiu MA. Paraventricular oxytocin neurons are involved in neural modulation of breathing. *J Appl Physiol* 92: 826–834, 2002.
- Martin-Body RL and Sinclair JD. Differences in respiratory patterns after acute and chronic pulmonary denervation. *Respir Physiol* 70: 205–219, 1987.
- McEwen B. Estrogen actions throughout the brain. *Recent Prog Horm Res* 57: 357–384, 2002.
- McEwen BS and Alves SE. Estrogen actions in the central nervous system. *Endocr Rev* 20: 279–307, 1999.
- Moore LG, McCullough RE, and Weil JV. Increased HVR in pregnancy: relationship to hormonal and metabolic changes. *J Appl Physiol* 62: 158–163, 1987.
- 34. **Pepe GJ and Albrecht ED.** Actions of placental and fetal adrenal steroid hormones in primate pregnancy. *Endocr Rev* 16: 608–648, 1995.
- Saaresranta T, Aittokallio T, Polo-Kantola P, Helenius H, and Polo O. Effect of medroxyprogesterone on inspiratory flow shapes during sleep in postmenopausal women. *Respir Physiol Neurobiol* 134: 131–143, 2003.
- Saaresranta T and Polo O. Hormones and breathing. Chest 122: 2165– 2182, 2002.
- Seralini G and Moslemi S. Aromatase inhibitors: past, present and future. Mol Cell Endocrinol 178: 117–131, 2001.
- Shahar E, Redline S, Young T, Boland LL, Baldwin CM, Nieto FJ, O'Connor GT, Rapoport DM, and Robbins JA. Hormone replacement therapy and sleep-disordered breathing. *Am J Respir Crit Care Med* 167: 1186–1192, 2003.

# R618

# PRENATAL ESTRADIOL AND DEVELOPMENT OF RESPIRATORY CONTROL

- Smith JC, Butera RJ, Koshiya N, Del Negro C, Wilson CG, and Johnson SM. Respiratory rhythm generation in neonatal and adult mammals: the hybrid pacemaker-network model. *Respir Physiol* 122: 131–147, 2000.
- Tatsumi K, Pickett CK, Jacoby CR, Weil JV, and Moore LG. Role of endogenous female hormones in hypoxic chemosensitivity. *J Appl Physiol* 83: 1706–1710, 1997.
- 41. Trotter A, Bokelmann B, Sorgo W, Bechinger-Kornhuber D, Heinemann H, Schmucker G, Oesterle M, Kohntop B, Brisch KH, and Pohlandt F. Follow-up examination at the age of 15 months of extremely preterm infants after postnatal estradiol and progesterone replacement. *J Clin Endocrinol Metab* 86: 601–603, 2001.
- 42. **Trotter A, Maier L, Grill HJ, Wudy SA, and Pohlandt F.** 17β-Estradiol and progesterone supplementation in extremely low-birth-weight infants. *Pediatr Res* 45: 489–493, 1999.
- 43. Trotter A, Maier L, and Pohlandt F. Management of the extremely preterm infant: is the replacement of estradiol and progesterone beneficial? *Pediatr Drugs* 3: 629–637, 2001.

- 44. Tsuruo Y, Ishimura K, Fujita H, and Osawa Y. Immunocytochemical localization of aromatase-containing neurons in the rat brain during preand postnatal development. *Cell Tissue Res* 278: 29–39, 1994.
- 45. Tsuruo Y, Ishimura K, Hayashi S, and Osawa Y. Immunohistochemical localization of estrogen receptors within aromatase-immunoreactive neurons in the fetal and neonatal rat brain. *Anat Embryol (Berl)* 193: 115–121, 1996.
- 46. Wang L, Andersson S, Warner M, and Gustafsson JA. Estrogen receptor (ER)β knockout mice reveal a role for ERβ in migration of cortical neurons in the developing brain. *Proc Natl Acad Sci USA* 100: 703–708, 2003.
- Young T, Finn L, Austin D, and Peterson A. Menopausal status and sleep-disordered breathing in the Wisconsin Sleep Cohort Study. *Am J Respir Crit Care Med* 167: 1181–1185, 2003.
- Zwain IH and Yen SS. Neurosteroidogenesis in astrocytes, oligodendrocytes, and neurons of cerebral cortex of rat brain. *Endocrinology* 140: 3843–3852, 1999.

